arXiv:1304.0014v1 [hep-th] 29 Mar 2013

ADM reduction of Einstein action and black hole entropy

I. Y. Park
Department of Applied Mathematics, Philander Smith College
Little Rock, AR 72223, USA
inyongpark05@gmail.com

Abstract

We carry out ADM reduction of the 4D Einstein-Hilbert action. The resulting theory is a scalar theory defined at a 3D hypersurface of the original black hole background, having an exponential potential. Once the the hypersurface is located at the Schwarzschild radius, the 3D theory is effectively reduced to a 2D Liouville type theory. We compute the partition function of the 2D theory, and show that its leading order entropy is proportional to the area of the horizon of the original black hole. The subleading terms come in logarithm/inverse powers of the area.

1 Introduction

Based on the previous works [1–3] (see [4–7] for related discussions), a new Kaluza-Klein scheme in which the bulk spacetime was viewed as a foliation of hypersurfaces along one of the coordinates, "r" was proposed in [8]. The scheme - called the ADM reduction - focuses on the collective dynamics of the hypersurfaces, which should capture aspects of the bulk physics. It is evident that the selected coordinate should play a special role, and this naturally suggests a description of dynamics where r plays the role of "time" coordinate.

Applied to a 5D supergravity theory, the ADM reduction led to the world-volume (one of the leaves) theory of the D3 brane [1] [8]. Here we apply the scheme to 4D pure Einstein gravity, and reduce it to a fixed point in the radial coordinate r. We show that the reduction leads to a 3D scalar theory description of the hypersurface being considered. Interestingly, the scalar theory takes the form of a 3D analogue of Liouville theory (and a 2D Liouville type theory subsequently) with an exponential potential.

In the big picture, the ADM scheme is to describe the physics of the 4D spacetime as fluctuations in the 3D and along the fourth direction. The first step in ADM reduction is ADM decomposition. The ADM reduction that we consider in this work is a relatively simple procedure that does not require the full machinery of ADM decomposition; below we will consider a reduction ansatz in the form of a block-diagonal decomposition,[1]

$$ds_4^2 \;=\; n^2 dr^2 + h_{\mu\nu} dx^\mu dx^\nu \tag{1}$$

Only the dynamics associated with fluctuations along r is allowed. (The rationale behind this reduction will be discussed below.) Then fluctuations will become dependent on the time and angle coordinates (t, θ, φ), and eventually, the system is rendered only of a scalar field in the fixed metric background $h_{\mu\nu}$. The scalar field - which we denote by $\phi(t, \theta, \varphi)$ below - is related to the radial component of the metric in a particular way that is presented in the next section.

In other words, the scalar theory describes a branch of the moduli space of the Einstein-Hilbert action, and should capture certain aspects of the 4D gravity theory. One may say that the particular branch of the moduli space of the original theory admits a "dual" description, which is the scalar theory in the present case. It seems, therefore, that the idea of "duality branch by branch [10]" is at work here too. In particular, it is plausible for the entropy of the BH system [11] [12] to be captured by the scalar theory. Discussions on microscopic origin of the black hole entropy can be found in [13–15].

---

[1] Compared with the standard ADM decomposition (see, e.g., [9] for a review), the shift vector is absent. Therefore, it should be viewed as a combination of ADM decomposition and Kaluza-Klein reduction in which the shift vector is set to zero.

One subtle issue is whether one should employ the finite temperature field theory technique to describe the dynamics of the hypersurface since the black hole has Hawking temperature. We believe that the zero-temperature field theory is the correct setup, and further comment on this issue in the conclusion.

The rest of the paper is organized as follows. In section 2, we implement the ADM reduction of 4D Einstein-Hilbert action. Although the actual reduction is carried out in the Lagrangian formulation, it is the Hamiltonian formulation that provides an intuitive and physical picture. After briefly reviewing the Hamiltonian formulation of "r-evolution", we demonstrate that the Schwarzschild metric can be obtained as a solution of this Hamiltonian formulation. When the hypersurface is taken to be the event horizon, only the spatial directions survive; the dimensions of the scalar theory are effectively reduced to two dimensions, and therefore, the theory becomes a Liouville type theory. Compared with the standard form of Liouville theory, the theory obtained has several inessential differences. Since our theory can be viewed as a minor variation of the standard Liouville theory, we will call the 2D theory just Liouville theory omitting the word "type". The computation of the entropy of the resulting 2D theory is undertaken in section 3 in which the area scaling of the entropy is established up to the issue of renormalization. The entropy obtained displays the same pattern that was observed in [16] (Entropy corrections were considered also in [17] [18], earlier). We conclude with summary and future directions. The appendix contains a discussion of entropy of a quantum field theory system at zero temperature.

2 ADM reduction of 4D Einstein action

In this section, we carry out ADM reduction of the 4D Einstein gravity. The ADM reduction that we consider in this work is a fairly straightforward procedure that does not require the full machinery of ADM decomposition but only its block-diagonal version. Roughly speaking, we leave arbitrary degrees of freedom to the (r, r)-component of the 4D metric $g_{\underline{\mu\nu}}$, writing it as $g_{rr} = g_{0rr} + \mathrm{g}_{\underset{\cdot}{r}r}$, and view $\mathrm{g}_{\underset{\cdot}{r}r}$ as the field that governs the dynamics of the hyersurface.

The actual reduction is carried out in the Lagrangian formulation for technical simplicity. However, it is the Hamiltonian formulation that provides an intuitive picture; we start by briefly reviewing the Hamiltonian formulation of "r-evolution" in section 2.1. We check in section 2.2 that the Schwarzschild metric can be obtained as a solution of the Hamiltonian formulation of r-evolution. In section 2.2, we carry out ADM reduction of the 4D Einstein-Hilbert action (this reduction might be somewhat similar in spirit to those of [19] [20]) and obtain Liouville field theory.

2.1 review of Hamiltonian formulation

We start with the Einstein-Hilbert action

$$S = \frac{1}{G} \int d^4x \sqrt{-g}\, R^{(4)} \tag{2}$$

The 4D coordinate $x^{\underline{\mu}}$ consists of the radial coordinate r and the 3D coordinate x^{μ}:

$$x^{\underline{\mu}} = r, x^{\mu} \quad \mu = 0, 1, 2 \tag{3}$$

Let us consider the following block-diagonal reduction ansatz for the solution[2],

$$ds_4^2 \;\; = \;\; n^2 dr^2 + h_{\mu\nu} dx^{\mu} dx^{\nu} \tag{4}$$

The 4D Einstein-Hilbert action now takes

$$\int d^4x \sqrt{-g^{(4)}} R^{(4)} = \int dr \int d^3x\, n\sqrt{-h} \left(\frac{1}{4n^2} (h^{\mu\nu} \partial_r h_{\mu\nu})^2 - \frac{1}{4n^2} (\partial_r h_{\mu\nu})^2 + R^{(3)} \right) \tag{5}$$

---

[2] It is more proper to include the shift vector; with it, the steps will be slightly modified (but with the same conclusion). However, it is crucial to omit the shift vector in section 2.2 where the reduction is carried out.

With the lagrangian given by

$$\mathcal{L} = n\sqrt{-h}\left(\frac{1}{4n^2}(h^{\mu\nu}\partial_r h_{\mu\nu})^2 - \frac{1}{4n^2}(\partial_r h_{\mu\nu})^2 + R^{(3)}\right) \tag{6}$$

the corresponding Hamiltonian is

$$\begin{aligned}\mathcal{H} &= p^{\mu\nu}\partial_r h_{\mu\nu} - \mathcal{L} \\ &= -n\sqrt{-h}\, R^{(3)} + \frac{n}{\sqrt{-h}}\Big(- p_{\mu\nu}^2 + \frac{1}{2}(p^\mu_\mu)^2\Big) \end{aligned} \tag{7}$$

where $p_{\mu\nu}$ is the canonical momentum, and ∇_μ is the 3D covariant derivative associated with the hypersurface under consideration. The field equations/constraints that follow from (7) are

$$\begin{aligned} -\dot{p}^{\mu\nu} &= n\sqrt{-h}\, G^{\mu\nu} + \frac{1}{2}\frac{n}{\sqrt{-h}}(p^{\rho\sigma}p_{\rho\sigma} - \frac{1}{2}p^2)h^{\mu\nu} - 2\frac{n}{\sqrt{-h}}(p^\mu_\rho p^{\nu\rho} - \frac{1}{2}pp^{\mu\nu}) \\ &\quad -\sqrt{-h}(\nabla^{(\nu}\nabla^{\mu)}n - h^{\mu\nu}\nabla^\rho\nabla_\rho n) \end{aligned} \tag{8}$$

$$-R^{(3)} + \frac{1}{(\sqrt{-h})^2}\Big(- p_{\mu\nu}^2 + \frac{1}{2}p^2\Big) = 0 \tag{9}$$

$$\dot{h}_{\mu\nu} = -2n\frac{1}{\sqrt{-h}}(p_{\mu\nu} - \frac{1}{2}h_{\mu\nu}p) \tag{10}$$

where $G^{\mu\nu} \equiv R^{\mu\nu} - \frac{1}{2}h^{\mu\nu}R$ and $p \equiv h_{\mu\nu}p^{\mu\nu}$. The constraint (9) shows that n is a moduli field: the on-shell Hamiltonian vanishes regardless of the value of n. The surface of a fixed metric still has the degrees of freedom associated with the moduli and, therefore, the entropy of the hypersurface with a fixed metric will naturally be associated with the dynamics of the moduli field n. Also note that

$$p_{\mu\nu} = \sqrt{-h}\Big(- \frac{1}{2n}\partial_r h_{\mu\nu} + h_{\mu\nu}\frac{1}{2n}h^{\rho\sigma}\partial_r h_{\rho\sigma}\Big) \tag{11}$$

and

$$\frac{1}{\sqrt{-h}}(-p_{\mu\nu}^2 + \frac{1}{2}p^2) = \sqrt{-h}\Big[- \frac{1}{4n^2}(\partial_r h_{\mu\nu})^2 + \frac{1}{4n^2}(h^{\rho\sigma}\partial_r h_{\rho\sigma})^2\Big] \tag{12}$$

The Hamiltonian formulation of r-evolution (7) admits, as it should, the Schwarzschild metric as a solution. One can check this explicitly as follows. Taking $n = n(r)$, the equation (8) becomes

$$-\frac{1}{n\sqrt{-h}}\partial_r p^{\mu\nu} = G^{\mu\nu} + \frac{1}{2}\frac{1}{(\sqrt{-h})^2}(p^2_{\rho\sigma} - \frac{1}{2}p^2)h^{\mu\nu} - \frac{2}{(\sqrt{-h})^2}(p^\mu_\rho p^{\nu\rho} - \frac{1}{2}pp^{\mu\nu}) \tag{13}$$

The trace part of the RHS of (13) implies

$$h_{\mu\nu}\partial_r p^{\mu\nu} = 0 \tag{14}$$

This and the non-trace part of (13) can easily be shown to lead to the solution of

$$n^2 = \left(1 - \frac{2GM}{r}\right)^{-1} \tag{15}$$

(The integration constant has been fixed in the usual manner.) Once the solution for $h_{\mu\nu}$ is obtained, $p_{\mu\nu}$ can be determined by (11). The result implies that the Schwarzschild metric

$$ds^2 = -\left(1 - \frac{2GM}{r}\right)dt^2 + \left(1 - \frac{2GM}{r}\right)^{-1} dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2) \tag{16}$$

is indeed a solution of the r-evolution Hamiltonian. For later use, we introduce the 3D metric solution, which we call "the 3D Schwarzschild metric",

$$\begin{aligned} ds^2_{3D} &= -\left(1 - \frac{2GM}{r}\right)dt^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2) \\ &= -\left(1 - \frac{2GM}{r}\right)dt^2 + \frac{4r^2}{(1+z\bar{z})^2}dz d\bar{z} \end{aligned} \tag{17}$$

in which the two-sphere part has been rewritten by introducing complex coordinates $z, \bar{z}$.

2.2 metric rescaling and reduction

One may consider the system given by (6), and in principle attempt to analyze the fluctuations around the Schwarzschild solution. However, the quantum level analysis would be complicated because of the non-canonical manner in which the moduli field n enters. In this section, we rescale the metric in order to make the system and its analysis more tractable. Towards the end of this section, we will note the effective reduction to 2D; it is that 2D theory whose entropy is computed in section 3.

Let us consider the Lagrangian (6),[3] and rescale the metric by

$$h_{\mu\nu} \to e^{\phi} h_{\mu\nu} \tag{18}$$

The Lagrangian now takes

$$\begin{aligned} \mathcal{L} \;=\;& n\sqrt{-h}\, e^{\frac{3}{2}\phi}\Big[\frac{3}{2n}\partial_r\phi + \frac{1}{2n}h^{\mu\nu}\partial_r h_{\mu\nu}\Big]^2 - n\sqrt{-h}\, e^{-\frac{1}{2}\phi}\Big[\frac{1}{2n}(\partial_r\phi)e^{\phi}h_{\mu\nu} + \frac{e^{\phi}}{2n}\partial_r h_{\mu\nu}\Big]^2 \\ & + n e^{\frac{1}{2}\phi}\sqrt{-h}\Big[R^{(3)} - 2\nabla_\mu\nabla^\mu\phi - \frac{1}{2}\nabla_\mu\phi\nabla^\mu\phi\Big] \end{aligned} \tag{19}$$

Once one imposes

$$e^{\phi} = \frac{1}{n^2} \tag{20}$$

one gets

$$\begin{aligned} =\;& \sqrt{-h}\, e^{2\phi}\Big(\frac{3}{2}\partial_r\phi + \frac{1}{2}h^{\mu\nu}\partial_r h_{\mu\nu}\Big)^2 - \sqrt{-h}\, e^{2\phi}\Big(\frac{1}{2}h_{\mu\nu}\partial_r\phi + \frac{1}{2}\partial_r h_{\mu\nu}\Big)^2 \\ & + \sqrt{-h}\Big[R^{(3)} - 2\nabla_\mu\nabla^\mu\phi - \frac{1}{2}\nabla_\mu\phi\nabla^\mu\phi\Big] \end{aligned} \tag{21}$$

Let us carry out the following reduction of the ϕ field

$$\phi(r,t,\theta,\varphi) \to \phi_0(r) + \phi(t,\theta,\varphi) \;\text{ with }\; e^{\phi_0} \equiv \frac{1}{n_0^2} \tag{22}$$

In this 3D setup, the functions that only have r-dependence may be treated as constants. The Lagrangian now takes

$$\begin{aligned} \mathcal{L}_{3D} \;=\;& \sqrt{-h}\, e^{2\phi}e^{2\phi_0}\left[\Big(\frac{3}{2}\partial_r\phi_0 + \frac{1}{2}h^{\mu\nu}\partial_r h_{\mu\nu}\Big)^2 - \Big(\frac{1}{2}h_{\mu\nu}\partial_r\phi_0 + \frac{1}{2}\partial_r h_{\mu\nu}\Big)^2\right] \\ & + \sqrt{-h}\Big[R^{(3)} - 2\nabla_\mu\nabla^\mu\phi - \frac{1}{2}\nabla_\mu\phi\nabla^\mu\phi\Big] \end{aligned} \tag{23}$$

---

[3] It turns out that once the shift vector is included, not only the ϕ-fluctuations but also the fluctuations of $h_{\mu\nu}$ must be included for consistency of the reduction.



Figure 1: reduction along r

One can show that the system admits (as it should) the following solution for the metric (with $\phi(t,\theta,\varphi)=0$),

$$ds^2_{3D} = h_{0\mu\nu}dx^\mu dx^\nu = -dt^2 + n_0^2 r^2(d\theta^2 + \sin^2\theta d\phi^2) \tag{24}$$

where n_0 denotes

$$n_0^2 = e^{-\phi_0} = \left(1 - \frac{2GM}{r}\right)^{-1} \tag{25}$$

(23) allows one to describe the physics of the surface of a fixed metric by analyzing the dynamics of ϕ.[4] This theory describes a branch of the moduli space of the hypersurface at an arbitrary $r \geq R_s$ location. Let us now place the hypersurface at the event horizon $r = R_S$ and consider fluctuations around the metric (24). Once the hypersurface is identified with the event horizon, $r = R_S$, the time-dependent configurations give contributions that oscillate rapidly in the path-integral, and therefore produce a suppressing factor due to the presence of the factor n_0. In the Minkowski case, this is due to Lebesgue lemma. The same is true for the Euclidean case that results from $t \to it$. In section 3, we will consider the 2D action that results from

---

[4] A more proper step would be integrating out the fluctuating metric around the fixed metric in the path integral. Once the metric background is chosen to be the Schwarzschild background, *the theory is in that curved background.* This with the work of [8] confirms once again that the proposal put forward in [21] was correct where it was conjectured that the holographic dual theory should be placed in the curved background. (See, e.g., [22] for further discussion.)

this Wick rotation. The theory reduces to 2D:

$$\begin{aligned}\mathcal{L}_{2D} &= \sqrt{-h}\, e^{2\phi} e^{2\phi_0} \left[\Big(\frac{3}{2}\partial_r\phi_0 + \frac{1}{2}h^{\mu\nu}\partial_r h_{\mu\nu}\Big)^2 - \Big(\frac{1}{2}h_{\mu\nu}\partial_r\phi_0 + \frac{1}{2}\partial_r h_{\mu\nu}\Big)^2\right] \\ &\quad + \sqrt{-h}\Big[R^{(3)} - \frac{1}{2}\nabla_a\phi\nabla^a\phi\Big] \qquad (26)\end{aligned}$$

where

$$a = 1, 2 \qquad (27)$$

and the total derivative part has been omitted. We gauge-fix the metric to

$$ds^2_{3D} = -dt^2 + h_{ab}dx^a dx^b \qquad (28)$$

and obtain

$$\begin{aligned}\mathcal{L}_{2D} &= \sqrt{-h}\, e^{2\phi} e^{2\phi_0} \left[\Big(\frac{3}{2}\partial_r\phi_0 + \frac{1}{2}h^{ab}\partial_r h_{ab}\Big)^2 - \Big(\frac{1}{2}\partial_r\phi_0\Big)^2 - \Big(\frac{1}{2}h_{ab}\partial_r\phi_0 + \frac{1}{2}\partial_r h_{ab}\Big)^2\right] \\ &\quad + \sqrt{-h}\Big[R^{(3)} - \frac{1}{2}\nabla_a\phi\nabla^a\phi\Big] \qquad (29)\end{aligned}$$

Note that only the 2D part of $R^{(3)}$ fluctuates. However, since 2D gravity is two-dimensional, it is not necessary to consider the fluctuations. (Namely, the gravity is now topological; the integrating out procedure mentioned in footnote 4 becomes irrelevant.) This system admits a round two-sphere metric for the metric part of the solution:

$$h_{0ab}dx^a dx^b = n_0^2 \, \frac{4r^2}{(1+z\bar{z})^2} dz d\bar{z} \qquad (30)$$

(There may be higher-genus solutions with the $\frac{4r^2}{(1+z\bar{z})^2}$ part replaced by higher genus surfaces; we will comment on them in the conclusion.) Let us substitute the solution (24) into the metric; it is straightforward to show

$$e^{2\phi_0} \left[\Big(\frac{3}{2}\partial_r\phi_0 + \frac{1}{2}h^{0\mu\nu}\partial_r h_{0\mu\nu}\Big)^2 - \Big(\frac{1}{2}h_{0\mu\nu}\partial_r\phi_0 + \frac{1}{2}\partial_r h_{0\mu\nu}\Big)^2\right] = \frac{2}{r^2}\Big(1 - \frac{2GM}{r}\Big)$$

and

$$R_0^{(3)} = \frac{2}{r^2}\Big(1 - \frac{2GM}{r}\Big)$$

r is to be set to $r = R_S$ after the n_0^2 factors are cancelled out. Omitting the "constant" $\sqrt{-h_0}\, R_0^{(3)}$ part, the Lagrangian can be rewritten as

$$S_{2D} \;=\; \int d^2x\Big(-\frac{1}{2}\partial_a\phi\partial^a\phi + 2\rho_0 e^{2\phi}\Big) \tag{31}$$

where

$$\rho_0 \equiv \frac{4}{(1+z\bar{z})^2} \tag{32}$$

Compared with the standard form of Liouville theory,[5] (31) is different from the standard form of Liouville theory in several ways. First, the kinetic term and the potential term have opposite signs. (We will come back to this point in the next section.) The linear term in ϕ is absent in (31).[6] As a matter of fact, a linear term appears once the factor ρ_0 in front of the $e^{2\phi}$ - which is another difference - is absorbed by shifting $\phi \to \phi + \tilde{\phi}_0$ with an appropriate function $\tilde{\phi}_0$:

$$\begin{aligned} S_{2D} \;&=\; \int d^2x\Big(-\partial_z\phi\partial_{\bar{z}}\phi + \frac{1}{2}\partial_z\phi\partial_{\bar{z}}\tilde{\phi}_0 + \frac{1}{2}\partial_z\tilde{\phi}_0\partial_{\bar{z}}\phi + e^{2\phi}\Big) \\ &=\; \int d^2x\Big(-\partial_z\phi\partial_{\bar{z}}\phi - \phi\partial_z\partial_{\bar{z}}\tilde{\phi}_0 + e^{2\phi}\Big) \end{aligned} \tag{33}$$

There is another contribution to the linear term. The field redefinition (18) should come with a determinant factor in the path integral over $h_{\mu\nu}$. The determinant factor brings the ϕ-linear term with an infinite coefficient (see, e.g., [29]). Therefore it is natural to expect that the linear term would arise through renormalization procedure even if it was absent in the beginning.

Although we will pursue the renormalization procedure elsewhere [30] (since it deserves thorough work in its own right), we sketch what needs to be established in that procedure. We find the issue of renormalization critical for several reasons. Firstly, the appearance of the linear term with an infinite coefficient that comes from the measure should be a strong indication of the need for renormalization at a fairly early stage of the analysis. Secondly, we will only consider the kinetic term and the exponential term in (33) in the next section. This linear term can be absorbed into the linear term coming from the measure. (One of the advantages of doing this is that the solution for the field equation is already known.) In terms of renormalization jargon, our setup amounts to starting with the bare action

---

[5] See, e.g., [23–26] for a review. Liouville theory appeared in the usual Kaluza-Klein context, e.g., in [27].

[6] Liouville thoery without the linear term was discussed for example in [28]. We thank Yu Nakayama for pointing out this reference.

$$S_{2D} \;=\; \int d^2x\Big(-\frac{1}{2}\partial_a\phi_B\partial^a\phi_B + Q_B R\phi_B + \mu_B e^{2b_B\phi_B}\Big) \tag{34}$$

and splitting it into the renormalized part and the counter terms[7]:

$$S_{2D} \;=\; \int d^2x\Big(-\frac{1}{2}\partial_a\phi_r\partial^a\phi_r + \mu_r e^{2b_r\phi_r}\Big) + \Delta S_{ctr} \tag{35}$$

where the counter-term part contains the linear term. Although we believe that such a renormalization program is very plausible, it must be explicitly established. We assume for now that this can be done. With this, our starting point for the entropy computation becomes

$$S_{2D} \;=\; \int d^2x\Big(-\partial_z\phi\partial_{\bar z}\phi + e^{2\phi}\Big) \tag{36}$$

where the subscript "r" is omitted and also in the following. Let us not turn to the entropy computation of this 2D theory.

3 Horizon-area scaling entropy from ϕ-field

In the previous section, we carried out the reduction along the radial (and time) direction(s) and obtained Liouville theory. It is actually (a version of) time-like Liouville theory due to the opposite signs of the kinetic term and the potential term.

In this section we compute the partition function and entropy. After reduction of the radial and time directions, an overall volume factor will come from

$$\int d^4x \Rightarrow V_{(t,r)}\int d^2x \quad , \quad V_{t,r} \equiv R_S^2\,\nu \tag{37}$$

where the volume associated with (t,r) directions has been defined in terms of the scale of the theory R_S and a dimensionless number ν. This step is crucial for the derivation of the area scaling as we will see shortly. The 2D action (36) becomes

---

[7] We later found out that a related discussion was given in [31].

$$S_{2D} = \int d^2x\Big(-\frac{R_S^2}{G}\partial_z\phi\partial_{\bar{z}}\phi + \frac{R_S^2}{G}\, e^{2\phi}\Big) \tag{38}$$

The constant ν has been has been absorbed to make the constant G bare with the prospect of renormalization. The G here denotes the renormalized coupling constant. By shifting ϕ by an appropriate constant, the $\frac{R_S^2}{G}$ in front of the potential $e^{2\phi}$ can be absorbed; one obtains

$$S_2 = \int d^2x\Big(-\frac{R_S^2}{G}\partial_z\phi\partial_{\bar{z}}\phi + e^{2\phi}\Big) \tag{39}$$

Let us use this form of action in the following to compute the entropy of the ϕ-system. Without this absorption of $\frac{R_S^2}{G}$ in front of $e^{2\phi}$, the perturbation series would diverge: suppose one uses (38) and insert several single-line vertices (i.e., the linear term), then the result will come with positive powers of $\frac{R_S^2}{G}$, and, therefore, the large-R_S perturbation will break down.

Let us find a solution of the system; the field equation that follows from (39) is

$$\partial_z\partial_{\bar{z}}\phi = -\frac{G}{R_S^2}e^{2\phi} \tag{40}$$

This admits the following solution as well known:

$$e^{2\phi_c} = \frac{R_S^2}{G}\frac{1}{(1+|z|^2)^2} \tag{41}$$

Next we carry out perturbation around this solution, obtaining the first two terms in the series for an illustration. Since we are interested in partition function, our task is simpler than computing general correlators of the interacting Liouville theory. Let us shift $\phi \to \phi_c + \phi$ where ϕ_c denotes the solution:

$$\begin{aligned} S_{2D} &= \int d^2x\Big(-\frac{R_S^2}{2G}\partial_a\phi_c\partial^a\phi_c - \frac{R_S^2}{2G}\partial_a\phi\partial^a\phi + e^{2\phi_c}(e^{2\phi}-2\phi)\Big) \\ &= -\frac{R_S^2}{G}\int d^2x\frac{1}{2}\partial_a\phi_c\partial^a\phi_c + \frac{R_S^2}{G}\int d^2z\left[-\frac{1}{2}\partial_a\phi\partial^a\phi + \frac{1}{(1+|z|^2)^2}(e^{2\phi}-2\phi)\right] \end{aligned} \tag{42}$$

The classical piece gives

$$-\frac{R_S^2}{G}\int d^2x\frac{1}{2}\partial_a\phi_c\partial^a\phi_c \quad = \quad -\frac{R_S^2}{2G}\left(\pi\ln\frac{R_S^2}{G}+const\right) \tag{43}$$

Although it may be possible to non-perturbatively evaluate the partition function by using the existing results in Liouville theory literature, we ponder the perturbative method for now because it reveals interesting patterns. The fluctuating part can be evaluated by the standard perturbation theory, which will produce a series in the "coupling constant" $\frac{G}{R_S^2}$. Consider the fluctuation part,

$$S_{quam}=\int d^2x\left(-\frac{1}{2}\frac{R_S^2}{G}\partial_a\phi\partial^a\phi+\frac{1}{(1+|z|^2)^2}(e^{2\phi}-2\phi)\right) \tag{44}$$

We may treat the exponential term of $(e^{2\phi}-2\phi)=2\phi^2+\cdots$ as vertices. The potential term has the "wrong sign" - we will comment on the potential significance of the sign in the conclusion. As far as the kinetic term is concerned, one choice of regularization seems to stick out: putting the system on a sphere of unit radius followed by zeta function regularization. This seems to be a natural choice given that the Lagrangian (29) admits a round sphere solution, and (42) has originated from the 2D action on two-sphere. Whether this regularization will also be effective for dealing with the vertices remains to be seen. The leading piece for the fluctuating part comes from the determinant of the kinetic part. The result depends on regularization method. (This seems to be a general feature of quantum entropy computations (see, e.g., [32]).) The 2D Laplacian is given by

$$\nabla^2 \quad = \quad \left[\frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\left(\sin\theta\frac{\partial}{\partial\theta}\right)+\frac{1}{\sin^2\theta}\frac{\partial^2}{\partial\varphi^2}\right] \tag{45}$$

and its eigenfunctions are the spherical harmonics:

$$\nabla^2 Y_{lm}=-l(l+1)Y_{lm} \tag{46}$$

One can use them to compute determinant of the propagator. Focusing on the R_S-dependent part, one finds

$$\left[\det\left(\frac{G}{R_S^2}\nabla^2\right)\right]^{-1/2}\sim e^{-\frac{1}{2}\mathrm{tr}\ln\frac{R_S^2}{G}} \tag{47}$$

Recording only the relevant part, the RHS can be rewritten as

$$e^{-\frac{1}{2}\ln\frac{R_S^2}{G}(\sum_{l=1}1)} = e^{\frac{1}{4}\ln\frac{R_S^2}{G}} \tag{48}$$

where the zero mode $l = 0$ has been omitted and zeta function regularization $\zeta(0) = -\frac{1}{2}$ has been used. Let us combine (43) and (48):

$$\exp\frac{R_s^2}{G}\left(-\frac{\pi}{2}\ln\frac{R_S^2}{G} + \frac{1}{4}\frac{G}{R_s^2}\ln\frac{R_S^2}{G} + \cdots\right) \tag{49}$$

where (...) denotes constant term and the terms of inverse powers of $\frac{R_s^2}{G}$ that come from the vertices. Unlike in the finite temperature field theory setup, the entropy cannot be defined as the temperature derivative of the free energy since the temperature does not explicitly enter our zero-temperature setup. Noting that the coupling constant (or the square of the coupling constant to be precise) plays the role of temperature, a derivative of the free energy with respect to the coupling constant (or its square) was taken to be the entropy analogue in [10] [32]. (See the appendix for a more detailed discussion.) In (39), $\frac{G}{R_S^2}$ plays the role of the coupling constant, and the entropy s can be defined in the same spirit as [10] [32]. The free energy analogue can be read off from (49):

$$F = -\frac{\pi}{2}\ln\frac{R_S^2}{G} + \frac{1}{4}\frac{G}{R_s^2}\ln\frac{R_S^2}{G} + \cdots \tag{50}$$

from which it follows

$$\begin{aligned} s &= \frac{\partial}{\partial(G/R_S^2)}\left[-\frac{\pi}{2}\ln\frac{R_S^2}{G} + \frac{1}{4}\frac{G}{R_s^2}\ln\frac{R_S^2}{G} + \cdots\right] \\ &= \frac{\pi}{2}\frac{R_S^2}{G} + \frac{1}{4}\ln\frac{R_S^2}{G} + \cdots \end{aligned} \tag{51}$$

where (...) contains constant term and terms of inverse powers of $\frac{R_S^2}{G}$. The coefficient of the logarithmic term is different from that of [16]. We comment on the possible reasons in the conclusion.

4 Conclusion

In this work, we have applied the ADM reduction scheme to the 4D Einstein-Hilbert action, and obtained the 3D system that describes the dynamics of a given hypersurface. When the location of the hypersurface is set at the Schwarzschild radius, the 3D theory is effectively reduced to Liouville theory.

There are several future directions. This work establishes (up to renormalization) that the BH entropy - which scales as area - comes from the 2D Liouville theory that is associated with the hypersurface. Careful study of normalization is needed to tie up some of the loose ends. The potential term in (44) is not bounded below; this should signify a certain instability. It will be interesting to see whether the instability could be associated with the evaporation of the black hole, and therefore the event horizon. Further work is needed to understand the origin of the coefficient of the logarithmic correction, which differs from the coefficient found in [16]. It should be possible to determine the coefficients of the terms of inverse powers of $\frac{R_S^2}{G}$ in (51). Another issue is temperature-related. We used the zero-temperature field theory setup even though the black hole has Hawking temperature. As wellknown, the black hole temperature was an induced effect in the original paper by Hawking. Therefore it is natural to expect that temperature should enter our zero-temperature setup (e.g., the 2D scalar theory) through certain dynamical (or possibly kinematical) effects. We have selected a hypersurface and did not consider “interactions” among the hypersurfaces. The role of the interactions of the neighboring hypersurfaces could be that they may induce finite temperature effects to the selected hypersurface. We plan to report on some of these issues in the near future.

Acknowledgements

I thank A. Nurmagambetov for useful discussions.

Appendix: Zero-T entropy of a QFT system

Conventionally, entropy of a quantum field theory system is considered in the finite-temperature setup. It is clear that in general a quantum field theory system at zero temperature should have finite entropy in general. We used a certain notion of zero-temperature entropy in [10] [32] and also in the present work. Although the notion is based on a close analogy with the finite temperature case, we discuss it here explicitly.

We limit our discussion to cases in which an Euclidean path-integral that takes the form of

$$\int d\Phi e^{-\frac{S(\Phi)}{\lambda}} \tag{A.1}$$

where Φ is the field that represents the degrees of freedom of the system and λ is the "coupling constant". Let us set this to

$$\int d\Phi e^{-\frac{S(\Phi)}{\lambda}} = e^{-\frac{F(\Phi,\lambda)}{\lambda}} \tag{A.2}$$

λ plays a role analogous to temperature in a finite temperature system, and F is an analogue of free energy [33]. Taking a derivative with respect to $1/\lambda$, one gets

$$\frac{\int d\Phi \; S e^{-\frac{S(\Phi)}{\lambda}}}{\int d\Phi e^{-\frac{S(\Phi)}{\lambda}}} = F + \frac{1}{\lambda}\frac{\partial F}{\partial(1/\lambda)} \tag{A.3}$$

The left-hand side is the average of the Hamiltonian, and the equation can be rewritten as

$$< H >= F - \lambda\frac{\partial F}{\partial\lambda} \tag{A.4}$$

where $< H >$ denotes the energy of the system. This implies that $\frac{\partial F}{\partial\lambda}$ should play a role analogous to the entropy of a finite-temperature system, s:

$$s \equiv \frac{\partial F}{\partial\lambda} \tag{A.5}$$

References

[1] M. Sato and A. Tsuchiya, “Born-Infeld action from supergravity,” Prog. Theor. Phys. **109**, 687 (2003) [hep-th/0211074].

[2] M. Sato and A. Tsuchiya, “Hamilton-Jacobi method and effective actions of D-brane and M-brane in supergravity,” Nucl. Phys. B **671**, 293 (2003) [hep-th/0305090].

[3] M. Sato and A. Tsuchiya, “M5-brane effective action as an on-shell action in supergravity,” JHEP **0411**, 067 (2004) [hep-th/0410261].

[4] J. Parry, D. S. Salopek and J. M. Stewart, “Solving the Hamilton-Jacobi equation for general relativity,” Phys. Rev. D **49**, 2872 (1994) [gr-qc/9310020].

[5] B. Darian, “Solving the Hamilton-Jacobi equation for gravitationally interacting electromagnetic and scalar fields,” Class. Quant. Grav. **15**, 143 (1998) [gr-qc/9707046].

[6] M. Fukuma, S. Matsuura and T. Sakai, “A Note on the Weyl anomaly in the holographic renormalization group,” Prog. Theor. Phys. **104**, 1089 (2000) [hep-th/0007062].

[7] T. Shiromizu, K. Koyama, S. Onda and T. Torii, “Can we live on a D-brane? Effective theory on a selfgravitating D-brane,” Phys. Rev. D **68**, 063506 (2003) [hep-th/0305253].

[8] E. Hatefi, A. J. Nurmagambetov and I. Y. Park, “ADM reduction of IIB on $H^{p,q}$ and dS braneworld,” arXiv:1210.3825 [hep-th].

[9] E. Poisson, “A relativist’s toolkit”, Cambridge university press (2004)

[10] E. Hatefi, A. J. Nurmagambetov and I. Y. Park, “N^3 entropy of M5 branes from dielectric effect,” Nucl. Phys. **B866**, 58-71 (2013) arXiv:1204.2711 [hep-th].

[11] J. D. Bekenstein, “Black holes and entropy,” Phys. Rev. D **7**, 2333 (1973).

[12] S. W. Hawking, “Black Holes and Thermodynamics,” Phys. Rev. D **13**, 191 (1976).

[13] G. 't Hooft, "On the Quantum Structure of a Black Hole," Nucl. Phys. B **256**, 727 (1985).

[14] L. Bombelli, R. K. Koul, J. Lee and R. D. Sorkin, "A Quantum Source of Entropy for Black Holes," Phys. Rev. D **34**, 373 (1986).

[15] M. Srednicki, "Entropy and area," Phys. Rev. Lett. **71**, 666 (1993) [hep-th/9303048].

[16] R. K. Kaul and P. Majumdar, "Logarithmic correction to the Bekenstein-Hawking entropy," Phys. Rev. Lett. **84**, 5255 (2000) [gr-qc/0002040].

[17] O. B. Zaslavsky, "Entropy of quantum fields for nonextreme black holes in the extreme limit," Phys. Rev. D **57**, 6265 (1998) [gr-qc/9708027].

[18] R. B. Mann, S. N. Solodukhin and , "Universality of quantum entropy for extreme black holes," Nucl. Phys. B **523**, 293 (1998) [hep-th/9709064].

[19] T. Biswas and W. Siegel, "Radial dimensional reduction: Anti-de Sitter theories from flat," JHEP **0207**, 005 (2002) [hep-th/0203115].

[20] A. Y. .Artsukevich and M. A. Vasiliev, "On Dimensional Degression in AdS(d)," Phys. Rev. D **79**, 045007 (2009) [arXiv:0810.2065 [hep-th]].

[21] I. Y. Park, "Fundamental versus solitonic description of D3-branes," Phys. Lett. B **468**, 213 (1999) [hep-th/9907142].

[22] I. Y. Park, "Open string engineering of D-brane geometry," JHEP **0808**, 026 (2008) [arXiv:0806.3330 [hep-th]].

[23] P. H. Ginsparg and G. W. Moore, "Lectures on 2-D gravity and 2-D string theory," [hep-th/9304011].

[24] V. Fateev, A. B. Zamolodchikov, A. B. Zamolodchikov and , "Boundary Liouville field theory. 1. Boundary state and boundary two point function," hep-th/0001012.

[25] J. Teschner, "Liouville theory revisited," Class. Quant. Grav. **18**, R153 (2001) [hep-th/0104158].

[26] Y. Nakayama, "Liouville field theory: A Decade after the revolution," Int. J. Mod. Phys. A **19**, 2771 (2004) [hep-th/0402009].

[27] H. Lu, C. N. Pope and K. W. Xu, "Liouville and Toda solutions of M theory," Mod. Phys. Lett. A **11**, 1785 (1996) [hep-th/9604058].

[28] M. Matone, "Uniformization theory and 2-D gravity. 1. Liouville action and intersection numbers," Int. J. Mod. Phys. A **10**, 289 (1995) [hep-th/9306150].

[29] A. A. Tseytlin, "Partition Function Of String Sigma Model On A Compact Two Space," Phys. Lett. B **223**, 165 (1989).

[30] Work in progress

[31] J. Distler, H. Kawai and , "Conformal Field Theory and 2D Quantum Gravity Or Who's Afraid of Joseph Liouville?," Nucl. Phys. B **321**, 509 (1989).

[32] I. Y. Park, "N^2 entropy of 4D N=4 SYM," arXiv:1208.1501 [hep-th].

[33] M. Peskin and D. Schroeder, "Quantum field theory", Addison-Wesley (1995)